

<b>County</b>	Prairie	<b>Upstream River Mile</b>	149.2
<b>Classification</b>	CM: Confined meandering	<b>Downstream River Mile</b>	137
<b>General Location</b>	To Terry Bridge	<b>Length</b>	12.20 mi (19.63 km)
<b>General Comments</b>	To Terry Bridge; confined		

## Narrative Summary

Reach D1 is located in Prairie County, and extends from just below the mouth of the Powder River to Terry. The reach is a 12.2 mile long Confined Meandering (CM) reach type, indicating that the river flows along a meandering course that is confined by older geologic units. Sandstones of the Fort Union Formation and younger erosion-resistant terraces confine the channel through the reach. Because of the geologic confinement, channel migration rates are low and the riparian corridor is notably thin or absent. There is one Fishing Access Site at the upper end of the reach at the Powder River confluence (Powder River Depot).

There are less than 1000 feet of bank armor in the reach; including about 550 feet of rock riprap and 140 feet of flow deflectors. The flow deflectors were all built between 2001 and 2011. During that timeframe there was a loss of 650 feet of rock riprap where it was protecting an old railroad bridge at RM 144.5. The bridge was built in 1907 for the railroad and now serves County Road 42.

Wolf Rapids is located on the apex of a large meander at RM 146. These rapids are formed from an exposed bedrock shelf that extends across the entire river.

Reach D1 has lost almost a mile of side channel length since 1950, but none of this loss has been associated with intentional blockages. There has been 126 acres of riparian recruitment into abandoned 1950s channels.

Land use is predominantly agricultural, and there has been 310 acres of land developed under pivot irrigation. There are two animal handling facilities just north of Terry that are adjacent to old swales. One dump site was mapped on the right bank of the river at RM 137.5R, about ¾ miles upstream from the Terry Bridge.

About 51 percent of the historic 5-year floodplain has become isolated, primarily due to flow alterations. The abandoned Milwaukee rail line embankment has been breached by river erosion in several locations on the south side of the river.

A total of four ice jams have been reported in the reach. One of these events was in February (1996), and three occurred in March (1993, 2009, and 2011). No damages were reported.

There are about 20 acres of mapped Russian olive in the reach.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The 2-year flood, which strongly influences overall channel form, has dropped by 22 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 4,850 cfs to 2,810 cfs with human development, a reduction of 42 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 6,940 cfs under unregulated conditions to 3,270 cfs under regulated conditions, a reduction of 53 percent.


CEA-Related observations in Reach D1 include:

- Breaching of abandoned Milwaukee Railroad line

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach D1 include:

- Dump site Practice at RM 137.5R
- Russian olive removal

## PHYSICAL FEATURES MAP (2011)



## HYDROLOGIC SUMMARY

Hydrologic data available for the Reach Narratives include data from representative gaging stations, modeling from the COE from the Big Horn river upstream, and modeling by the USGS for the Big Horn River to the Missouri River confluence. Gaging stations that best represent the watershed area within any reach are used to describe the flood history within the reach. Hydrology modeling results generated for all reaches provides unregulated and regulated flow values. Seasonal and annual flow duration data generated by the USGS are available for reaches C10 through D13.

### Gage Representation (Gage-Based): Sidney

#### Flood History

Year	Date	Flow on Date	Return Interval	Gage No	Downstream Gage	Upstream Gage
1978	May 23	111,000	10-25 yr	6329500	6309000	
1912	Mar 29	114,000	10-25 yr	Location	Sidney	Miles City
1944	Jun 21	120,000	10-25 yr	Period of Record	1911-2015	1929-2015
2011	May 24	124,000	10-25 yr	Distance To (miles)	106.2	34.8
1918	Jun 20	126,000	25-50 yr			
1943	Mar 29	132,000	25-50 yr			
1923	Oct 3	134,000	25-50 yr			
1952	Mar 31	138,000	25-50 yr			
1921	Jun 21	159,000	100-yr			

#### Discharge

	1.01 Yr	2 Yr	5 Yr	10 Yr	50 Yr	100 Yr	500 Yr	7Q10 Summer	95% Sum. Duration
Unregulated		68,200	87,300	99,900	128,000	140,000	169,000	4,850	6,940
Regulated		53,000	71,700	83,600	109,000	119,000	142,000	2,810	3,280
% Change		-22.29%	-17.87%	-16.32%	-14.84%	-15.00%	-15.98%	-42.06%	-52.74%

#### Flow Duration

Streamflow, in ft<sup>3</sup>/s, which was equaled or exceeded for indicated percent of time

Note that these statistics are only available from Reach C10 downstream. See the USGS report for detailed information.

Season		5%	50%	95%
Spring	Unregulated	66,400	24,800	6,810
	Regulated	51,200	14,800	4,970
	% Change	-23%	-40%	-27%
Summer	Unregulated	46,600	14,700	6,940
	Regulated	34,800	8,850	3,280
	% Change	-25%	-40%	-53%
Fall	Unregulated	9,690	5,920	2,090
	Regulated	11,100	7,380	3,610
	% Change	15%	25%	73%
Winter	Unregulated	14,100	5,300	2,100
	Regulated	14,700	6,450	3,410
	% Change	4%	22%	62%
Annual	Unregulated	49,200	8,790	2,830
	Regulated	36,700	7,940	3,670
	% Change	-25%	-10%	30%

## AERIAL PHOTOGRAPHY

A variety of aerial photographic sources provide the basis for much of the Cumulative Effects Assessment analysis. The table below lists the air photos compiled for the reach and the associated discharge at the most representative USGS gaging station.

	Source	Acquisition Date	Type	Scale	Gage	Discharge
1950	USGS-EROS	26-Aug-49	B/W	1:14,800	6329500	2750
1976	USCOE	9-Oct-76	B/W	1:24,000	6329500	9580
1995	USGS DOQQ	1995??	B/W		6329500	
2001	NRCS	August 2-8, 2001	CIR	1:24,000	6329500	4000
2005	NAIP	08/05/2005	color	1-meter pixels	6329500	4170
2005	NAIP	08/04/2005	color	1-meter pixels	6329500	4350
2007	Woolpert	10/15/2007 - 11/2/0007	Color		6329500	
2009	NAIP	8/11/2009	Color	1-meter pixels	6329500	13000
2011	USCOE	October 2012	color	1-ft pixel	6329500	9030
2011	NAIP	7/25/2011	Color	1-meter pixels	6329500	41100
2011	NAIP	7/16/2011	Color	1-meter pixels	6329500	60100
2013	NAIP	07/24/2013	color	1-meter pixels	6329500	
2013	NAIP	07/19/2013	color	1-meter pixels	6329500	

## PHYSICAL FEATURES

Several efforts to capture the types and extents of physical features in the corridor have been generated by the CEA study. The 2001 Physical Features Inventory was performed through helicopter/video Rapid Aerial Assessment by the NRCS (NRCS, 2001) and did not include Park County. This inventory includes point and linear features that represent bank armor, irrigation structures, transportation encroachments, and areas of accelerated erosion. Bank armor mapped in the 2001 inventory only reflects features on the active channel margin, and thus excludes off-channel features on historic side channels. Some floodplain restriction features such as dikes and levees in the 2001 Physical Features Inventory may extend well beyond the active channel. In 2013, the 2001 inventory was revised to include Park County. At that time, some attribute inconsistencies in the original data were addressed. This dataset was then updated to reflect conditions in the 2011 NAIP imagery.

For Stillwater, Yellowstone and Dawson Counties, a Physical Features Timeline was generated that includes additional mapping based on aerial photography and assigns approximate dates of feature construction based on observed presence/absence in historic imagery between the 1950s and 2005 (DTM and AGI, 2008). The Physical Features Timeline contains features that were not mapped in the 2001 inventory (e.g. bank armor abandoned in floodplain areas by 2001). As such the total bank armor extent in the 2005 data is commonly greater than that identified in 2001 or 2013.

Note: As the goal for each physical features mapping effort were different, with differing mapping extents, there will be discrepancies between total feature lengths (e.g. length of rock riprap) in each data set.

### 2001 and 2011 Physical Features Bankline Inventories

Feature Class	Feature Type	2001 Length (ft)	% of Bankline	2011 Length (ft)	% of Bankline	2001-2011 Change
Stream Stabilization						
	Rock RipRap	1,196	0.9%	545	0.4%	-651
	Flow Deflectors	0	0.0%	113	0.1%	113
	Between Flow Deflectors	0	0.0%	130	0.1%	130
	<b>Feature Type Totals</b>	<b>1,196</b>	<b>0.9%</b>	<b>787</b>	<b>0.6%</b>	<b>-409</b>
Other In Channel						
	Bedrock Outcrop	1,429	1.1%	1,429	1.1%	0
	<b>Feature Type Totals</b>	<b>1,429</b>	<b>1.1%</b>	<b>1,429</b>	<b>1.1%</b>	<b>0</b>
Floodplain Control						
	Floodplain Dike/Levee	4,290	3.4%	4,290	3.4%	0
	<b>Feature Type Totals</b>	<b>4,290</b>	<b>3.4%</b>	<b>4,290</b>	<b>3.4%</b>	<b>0</b>
	<b>Reach Totals</b>	<b>6,914</b>	<b>5.4%</b>	<b>6,506</b>	<b>5.1%</b>	<b>-409</b>


### Intent of Bank Protection: 2001

The 2001 bank protection features were assessed for the 'intent' of what they protect.

Feature Type	Irrigated	Non-Irrig.	Ag. Infrastr.	Road	Interstate	Railroad	Urban	Exurban
Rock RipRap	0	0	0	1,197	0	0	0	0
<b>Totals</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1,197</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>

## ICE JAMS

Ice jam data were obtained from the National Ice Jam Database maintained by the Ice Engineering Group at Army Corps of Engineers Cold Regions Research and Engineering Laboratory (<https://rsgis.crrel.usace.army.mil/icejam/>). From this database, Yellowstone River ice jams are summarized by reach in the Yellowstone River Historic Events Timeline (DTM and AGI, 2008b). The basic information for each ice jam is presented as a list of events. The graph represents the number of database entries for a reach. Note that a single jam event may have multiple entries.



Jam Date	Jam Type	River Mile	Damages
3/26/1993	Break-up	137	?
2/9/1996	Break-up	137	?
3/3/2009	Break-up		None known
3/14/2011	Break-up		

## GEOMORPHIC

The geomorphology data presented below consist of measured changes in Braiding Parameter since 1950 and blocked side channels. Braiding parameter is a measure of the total length of side channels relative to that of the main channel. The braiding parameter is calculated as the sum of anabranching and primary channel lengths divided by the primary channel length. Secondary channels within the bankfull margins are a function of flow stage and hence were not included in the braiding parameter calculation. If a reach has a braiding parameter of 3, then the total bankfull channel length is three times that of the main channel. The mean braiding parameter measured for all 88 reaches is 1.8.

Blocked side channels that were either plugged with a small dike or cutoff by larger features such as a levee or road prism were identified for the pre and post-1950s eras.

Additional geomorphic parameters are discussed in more detail in the study report and appendices.

### Braiding (Bankfull)

	Primary Chan. Length (ft)	Anab. Ch. Length (ft)	Bankfull Braiding Parameter	% Change in Braiding
1950	65,951	16,560	1.25	1950 to 1976: 2.31%
1976	63,797	17,862	1.28	1976 to 1995: -6.65%
1995	63,973	12,462	1.19	1995 to 2001: 0.01%
2001	63,919	12,462	1.19	1950 to 2001: -4.49%
Change 1950 - 2001	-2,032	-4,098	-0.06	

Length of Side Channels Blocked	Pre-1950s (ft)	Post-1950s (ft)
	0	0

## HYDRAULICS

Available hydraulic information includes county-based HEC-RAS modeling efforts by the Army Corps of Engineers with the exclusion of Park County. Floodplain modeling was performed for four conditions representing a developed and undeveloped floodplain, and unregulated and regulated flows for the 1.5, 2, 5, 10, 20, 50, 100, 200, and 500-year events. Park County has limited FEMA hydraulic modeling and was not included in the analysis.

The results of HEC-RAS modeling for the 5 and 100-year flood events were assessed to compare the extents of inundated area for the pristine (undeveloped floodplain, unregulated flows) and developed (developed floodplain, regulated flows) conditions. The data sets provided for each flow condition were unioned in the GIS to identify areas where the inundated extent differed. These areas of human-caused floodplain isolation due to either flow alterations or physical features such as levees. For the 100-year flood event, isolated areas greater than 5 acres were attributed with the interpreted reason for isolation (railroad, levee, etc.). The resulting values are presented as acres and percent of the pristine floodplain that has been isolated. The pristine floodplain is defined as the total floodplain footprint minus the area of the mapped 2001 bankfull channel (mapped islands were included in the floodplain area).

### Floodplain Isolation

	100-Year		5-Year	
	Isolated Acres	% of Floodplain	Isolated Acres	% of Floodplain
Non-Structural (hydrology, geomorphic, etc.)	0	0.0%		
Agriculture (generally relates to field boundaries)	0	0.0%		
Agriculture (isolated by canal or large ditch)	0	0.0%		
Levee/Riprap (protecting agricultural lands)	15	3.3%		
Levee/Riprap (protecting urban, industrial, etc.)	0	0.0%		
Railroad	0	0.0%		
Abandoned Railroad	0	0.0%		
Transportation (Interstate and other roads)	0	0.0%		
<b>Total Not Isolated (Ac)</b>	<b>440</b>		<b>1321</b>	
<b>Total Floodplain Area (Ac)</b>	<b>455</b>		<b>1417</b>	
<b>Total Isolated (Ac)</b>	<b>15</b>	<b>3.3%</b>	<b>95</b>	<b>51.3%</b>

The 5-year floodplain is a good allegory for the extent of the riparian zone. Thus, irrigated areas within the 5-year floodplain tend to represent riparian zones that have been converted to agriculture and may result in additional bank protection to protect the agricultural production and irrigation infrastructure.

	Flood	Sprinkler	Pivot	Total
Irrigated Acres within the 5 Year Flooplain:	0	0	0	0

## CHANNEL MIGRATION ZONE

A series of Channel Migration Maps were developed for the Yellowstone River from Gardiner to its mouth in McKenzie County, North Dakota (Thatcher, Swindell, and Boyd, 2009). These maps and their accompanying report can be accessed from the YRCDC Website. The channel migration zone (CMZ) developed for the Yellowstone River is defined as a composite area made up of the existing channel, the historic channel since 1950 (Historic Migration Zone, or HMZ), and an Erosion Buffer that encompasses areas prone to channel erosion over the next 100 years. Areas within this CMZ that have been isolated by constructed features such as armor or floodplain dikes are attributed as “Restricted Migration Areas” (RMA). Beyond the CMZ boundaries, outlying areas that pose risks of channel avulsion are identified as “Avulsion Potential Zones”.

Mean 50-Yr Migration Distance (ft)	Erosion Buffer (ft)	Total CMZ Acreage	Restricted CMZ Acreage	% Restricted Migration Area	Total AHZ Acreage	Restricted AHZ Acreage	% Restricted Avulsion Area
120	241	1,889	12	1%	0	0	0%

### 2011 Restricted Migration Area Summary

Note that these data reflect the observed conditions in the 2011 aerial photography (NAIP for Park and Sweet Grass Counties, COE for the rest of the river).

Reason for Restriction	Land Use Protected	RMA Acres	Percent of CMZ
Road/Railroad Prism			
	Public Road	12	0.6%
<b>Totals</b>		<b>12</b>	<b>0.6%</b>

### Land Uses within the CMZ (Acres)

Flood Irrigation	Sprinkler Irrigation	Pivot Irrigation	Urban/ExUrban	Transportation
21.7	0.0	2.0	0.1	5.1



## LAND USE

Land uses were mapped from aerial photography Gardiner to the confluence of the Missouri River in North Dakota for four time periods: 1950s, 1976, 2001, and 2011. Mapping was performed at approximately 1:6,000 to ensure consistent mapping across all data sets. Typically, if a feature could not be easily mapped at the target mapping scale, it was not separated out from the adjacent land use.

A four-tiered system was used to allow analysis at a variety of levels. Tier 1 breaks land use into Agricultural and Non-Agricultural uses. Tier two subdivided uses into productive Agricultural Land and Infrastructure for the Agricultural land, and Urban, Exurban and Transportation categories for the Non-Agricultural land. Tier three further breaks down land uses into more refined categories such as Irrigated or Non-Irrigated and Residential, Commercial, or Industrial. Finally, Tier 4 focuses primarily on the productive agricultural lands, identifying the type of irrigation (Pivot, Sprinkler or Flood).

### Land Use Timeline - Tiers 2 and 3

Feature Class	Feature Type	Acres				% of Reach Area			
		1950	1976	2001	2011	1950	1976	2001	2011
Agricultural Infrastructure									
	Canal	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Agricultural Roads	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Other Infrastructure	7	31	64	57	0.1%	0.4%	0.8%	0.7%
	<b>Totals</b>	<b>7</b>	<b>31</b>	<b>64</b>	<b>57</b>	<b>0.1%</b>	<b>0.4%</b>	<b>0.8%</b>	<b>0.7%</b>
Agricultural Land									
	Non-Irrigated	5,846	5,654	5,499	5,383	73.8%	71.4%	69.4%	67.9%
	Irrigated	682	866	1,048	1,157	8.6%	10.9%	13.2%	14.6%
	<b>Totals</b>	<b>6,529</b>	<b>6,519</b>	<b>6,547</b>	<b>6,540</b>	<b>82.4%</b>	<b>82.3%</b>	<b>82.6%</b>	<b>82.5%</b>
Channel									
	Channel	1,283	1,257	1,221	1,251	16.2%	15.9%	15.4%	15.8%
	<b>Totals</b>	<b>1,283</b>	<b>1,257</b>	<b>1,221</b>	<b>1,251</b>	<b>16.2%</b>	<b>15.9%</b>	<b>15.4%</b>	<b>15.8%</b>
ExUrban									
	ExUrban Other	0	16	16	16	0.0%	0.2%	0.2%	0.2%
	ExUrban Undeveloped	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Industrial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Commercial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	ExUrban Residential	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	<b>Totals</b>	<b>0</b>	<b>16</b>	<b>16</b>	<b>16</b>	<b>0.0%</b>	<b>0.2%</b>	<b>0.2%</b>	<b>0.2%</b>
Transportation									
	Public Road	31	25	42	27	0.4%	0.3%	0.5%	0.3%
	Interstate	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Railroad	72	74	32	32	0.9%	0.9%	0.4%	0.4%
	<b>Totals</b>	<b>104</b>	<b>99</b>	<b>74</b>	<b>59</b>	<b>1.3%</b>	<b>1.2%</b>	<b>0.9%</b>	<b>0.7%</b>
Urban									
	Urban Other	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Residential	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Commercial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Undeveloped	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	Urban Industrial	0	0	0	0	0.0%	0.0%	0.0%	0.0%
	<b>Totals</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>	<b>0.0%</b>

### Land Use Timeline - Tiers 3 and 4

Feature Class	Feature Type	Acres				% of Reach Area				Change Between Years (% of Agricultural Land)			
		1950	1976	2001	2011	1950	1976	2001	2011	'50-76	'76-01	'01-11	'50-11
Irrigated													
	Sprinkler	0	0	0	0	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
	Pivot	0	0	202	310	0.0%	0.0%	3.1%	4.7%	0.0%	3.1%	1.7%	4.7%
	Flood	682	866	846	846	10.5%	13.3%	12.9%	12.9%	2.8%	-0.4%	0.0%	2.5%
	<b>Totals</b>	<b>682</b>	<b>866</b>	<b>1,048</b>	<b>1,157</b>	<b>10.5%</b>	<b>13.3%</b>	<b>16.0%</b>	<b>17.7%</b>	<b>2.8%</b>	<b>2.7%</b>	<b>1.7%</b>	<b>7.2%</b>

Non-Irrigated

Multi-Use	5,405	5,129	5,205	5,212	82.8%	78.7%	79.5%	79.7%	-4.1%	0.8%	0.2%	-3.1%
Hay/Pasture	441	524	294	171	6.8%	8.0%	4.5%	2.6%	1.3%	-3.5%	-1.9%	-4.1%
<b>Totals</b>	<b>5,846</b>	<b>5,654</b>	<b>5,499</b>	<b>5,383</b>	<b>89.5%</b>	<b>86.7%</b>	<b>84.0%</b>	<b>82.3%</b>	<b>-2.8%</b>	<b>-2.7%</b>	<b>-1.7%</b>	<b>-7.2%</b>

## RIPARIAN

Riparian mapping data are derived from the Yellowstone River Riparian Vegetation Mapping study (DTM/AGI 2008). This study coarsely mapped the riparian vegetation communities using 1950's, 1976-1977, and 2001 aerial imagery in a GIS environment. The polygons are digitized at a scale of approximately 1:7,500, with a minimum mapping unit of approximately 10 acres. The goal of the delineation was to capture areas of similar vegetation structure as they appeared on the aerial imagery, while maintaining a consistent scale.

The "Riparian Turnover" values quantify the total area within the active channel area that converted from either woody vegetation to open bar or water, or from open bar or water to woody vegetation. A comparison of these values allows some consideration of overall riparian encroachment into the river corridor from 1950 to 2001.

### Riparian Mapping

Statistic	Shrub (Acres)			Closed Timber (Acres)			Open Timber (Acres)		
	1950	1976	2001	1950	1976	2001	1950	1976	2001
Min	1.6	0.9	0.2	3.3	1.7	0.7	2.9	4.8	3.9
Max	17.6	33.6	21.6	32.4	19.7	23.1	13.0	23.3	17.3
Average	5.0	5.1	5.0	14.7	6.4	5.5	6.6	13.4	11.1
Sum	69.7	97.7	44.9	88.1	44.5	43.8	39.8	80.5	66.3

### Riparian Turnover

Conversion of riparian areas to channel, or from channel to riparian between the 1950's and 2001 data set.

Riparian to Channel (acres) 27.1

Channel to Riparian (acres) 34.2

**Riparian Encroachment (acres) 7.2**

### Riparian Recruitment

Creation of riparian areas between 1950s and 2001.	1950s Channel Mapped as 2011 Riparian (Ac)	126.0
	1950s Floodplain Mapped as 2011 Channel (Ac)	7.6
	<b>Total Recruitment (1950s to 2011)(Ac)</b>	<b>133.6</b>

## WETLANDS

Wetland areas were mapped to National Wetland Inventory standards by the Montana Natural Heritage Program. Palustrine wetlands within the mapped 100-year inundation boundary were extracted and summarized into four categories: Riverine (Unconsolidated Bottom - UB, Aquatic Bed - AB, and Unconsolidated Shore - US), Emergent - EM, Scrub-Shrub - SS, and Forested - FO.

	Riverine	Emergent	Scrub/Shrub	Forested	Total
<b>Mapped Acres</b>	27.0	18.0	0.0	0.0	<b>45.0</b>
<b>Acres/Valley Mile</b>	3.0	2.0	0.0	0.0	

## RUSSIAN OLIVE

Russian olive is considered an invasive species and its presence in the Yellowstone River corridor is fairly recent. As such, its spread can be used as a general indicator of invasive plants within the corridor. It has the added benefit of being easily identified in multi-spectral aerial photography, making it possible to inventory large areas using remote techniques.

In 2011, Natural Resources Conservation Service (NRCS) in Bozeman, MT conducted an inventory of Russian olive locations in the Yellowstone River watershed. This study utilized the Feature Analyst extension within ArcGIS to interpret multi-spectral 2008 NAIP imagery for the presence of Russian olive. The resulting analysis was converted from raster format to a polygon ESRI shape file for distribution and further analysis within a GIS environment.

This work scope was tasked with integrating the resulting Russian olive inventory into the Yellowstone River Conservation Districts Council (YRDC) Cumulative Effects Assessment (CEA) GIS and associated reach-based database. Additionally, analysis of Russian olive within the corridor was conducted to characterize its distribution in throughout the corridor and its association with other corridor data sets.

	Floodplain Area (Ac)	% of Floodplain	Other Area (Ac)	Inside RMA (Ac)	Inside '50s Channel (Ac)	Inside 50s Island (Ac)
<b>Russian Olive in Reach</b>	19.88	1.36%	40.51	0.07	6.50	0.37

## FISHERIES SUMMARY

Fisheries data available for the Reach Narratives include low-flow and high-flow habitat mapping of 2001 conditions for 406 miles of river, extending from the mouth upstream to a point approximately 8 miles upstream of Park City. Habitat mapping was performed remotely on the 2001 CIR aerial photography utilizing habitat classifications developed by Montana Fish, Wildlife, and Parks (DTM 2009). Historic habitat mapping using the 1950's imagery is limited to Reach B1 (high-flow) and D9 (low and high-flow).

Fisheries field sampling data have been provided by Ann Marie Reinhold (MSU). In this study, the Yellowstone River from Park City to Sidney was divided into five segments. Within each segment, fish were sampled in reaches modified by riprap ("treatment reaches") and relatively unmodified reaches ("control reaches"). Fish sampling was conducted during summer and autumn of 2009, 2010, and 2011. Boat electrofishing, trammel nets, mini-fyke nets and bag seines were used to collect data from river bends.

Fish presence data is only presented for those reaches that were sampled.

The Low Flow Habitat Mapping followed schema developed by Montana Fish Wildlife and Parks to identify key habitat units for certain aquatic species.

### Low Flow Fisheries Habitat Mapping

Habitat	2001 (Acres)		
	Bankfull	Low Flow	% of Low Flow
Scour Pool	90.2	81.8	6.4%
Rip Rap Bottom	78.2	48.0	3.8%
Bluff Pool	235.1	183.6	14.4%
Terrace Pool	350.6	212.5	16.7%
Secondary Channel	63.2	40.6	3.2%
Secondary Channel (Seasonal)	29.4	49.1	3.9%
Channel Crossover	373.3	299.2	23.5%
Point Bar		105.0	8.3%
Side Bar		117.9	9.3%
Mid-channel Bar		66.6	5.2%
Island	50.1	23.1	1.8%
Dry Channel		41.5	3.3%
Confluence Area		2.7	0.2%

## AVIAN

Birds were sampled in 2006 and 2007 by Danielle Jones of Montana State University. Point count methods were used at 304 randomly chosen sites in 21 braided or anabranching reaches. Each site was visited multiple times within a season, and sites were visited in both years. Birds were sampled in grassland, shrubland, and cottonwood forest habitats. Additional bird data was collected by Amy Cilimburg of Montana Audubon in summer 2012. High priority areas for data collection were identified with the assistance of the YRCDC Technical Advisory Committee. The Audubon methodology recorded data for a wider variety of bird species relative to the MSU study, including raptors and waterfowl.

## CULTURAL INVENTORY SUMMARY

The Yellowstone River Cultural Inventory - 2006 documents the variety and intensity of different perspectives and values held by people who share the Yellowstone River. Between May and November of 2006, a total of 313 individuals participated in the study. They represented agricultural, civic, recreational, or residential interest groups. Also, individuals from the Crow and the Northern Cheyenne tribes were included.

There are three particular goals associated with the investigation. The first goal is to document how the people of the Yellowstone River describe the physical character of the river and how they think the physical processes, such as floods and erosion, should be managed. Within this goal, efforts have been made to document participants' views regarding the many different bank stabilization techniques employed by landowners. The second goal is to document the degree to which the riparian zone associated with the river is recognized and valued by the participants. The third goal is to document concerns regarding the management of the river's resources. Special attention is given to the ways in which residents from diverse geographical settings and diverse interest groups view river management and uses. The results illustrate the commonalities of thought and the complexities of concerns expressed by those who share the resources of the Yellowstone River.

### Summary of Cultural Views in Region D

A review of the interview data for the segment, Missouri River to Powder River, suggests that people in this area engage in four primary discussions when asked about the Yellowstone River. First, the notion of Eastern Montana is not simply a geographic reference. It is a defining concept that captures the agricultural roots and the cultural values of the people living in the study segment, and the river is an essential element within their notion of Eastern Montana. Second, the river is discussed as a wholesome recreational outlet. However, shifting landownership is noted as an important change in the recreational context. Third, even though agricultural practices are viewed as the mainstay of the local economies, many participants discuss the long-term economic viability of their communities as a concern. Industrial and residential developments along the river's edge are seemingly remote possibilities and are generally discussed with references to flood plain restrictions and the stability of nearby dikes. Finally, discussions of managing the river are limited, but a variety of opinions are offered regarding bank erosion and stabilization techniques.