

Yellowstone River Reach Narratives

Reach C16

County	Custer	Upstream River Mile	192.3
Classification	PCM/I: Partially confined meandering/islands	Downstream River Mile	185
General Location	to Miles City	Length	7.30 mi (11.75 km)

Narrative Summary

Reach C16 is 7.32 miles long and is located just upstream of Miles City. The downstream limit of the reach is the mouth of the Tongue River at RM 185. The reach is characterized by a dominant main thread that shows a distinct meandering pattern, with several islands persisting where meander bends have historically cut off. The river intermittently flows along the valley wall. As a result it is classified as Partially Confined Meandering with Islands (PCM/I).

As of 2011 there were about two miles of armor protecting 14 percent of the total bankline in Reach C16, including 7,000 feet of rock riprap, 2,200 feet of concrete riprap, and 1,550 feet of flow deflectors. All of the concrete armor is protecting urban areas around the water treatment plant in Miles City. The flow deflectors protect non-irrigated agricultural land, and the rock riprap is protecting agricultural land (irrigated and non-irrigated), roads, and the rail line. A ~550 foot-long stretch of armor at RM 190.5R has been flanked since 2001, and erosion behind the armor now threatens a road; the river has locally eroded into the road embankment. There were also several miles of transportation encroachments and floodplain levees mapped in the reach.

About 13 percent (308 acres) of the 100-year floodplain has become isolated from the river in Reach C16, meaning it is no longer inundated at what was historically a 100-year flood event. Isolation can be due to flow changes and/or physical features that block overflows from reaching floodplain areas. Most of the 100-year floodplain isolation (185 acres) is due to the active rail line. Isolation of the 5-year floodplain has been even more substantial, with 62 percent (721 acres) of the historic 5-year floodplain no longer inundated at what was historically a 5-year flood event.

Three ice jams have been reported in the reach, including February of 2011, and March of 2003 and 2012. No damages were recorded in the ice jam database.

At RM 186.6 a steel trestle bridge built for the now abandoned Milwaukee Railroad crosses the river where it is about 1,000 feet wide. There are several very large barbs on the right bank of the river upstream of the bridge that extend about 100 feet off of the bank, and there is riprap directly under the structure.

About 210 acres which represents 9 percent of the total CMZ have become restricted by physical features. Areas that have become restricted to channel migration include the water treatment plant just upstream of the mouth of the Tongue River, behind the railroad grade at RM 191.5, and locally behind stretches of bank armor protecting irrigated and non-irrigated fields.

Mapped land uses in Reach C16 range from agricultural to urban to transportation infrastructure. The total acreage of flood irrigated land in the reach has dropped from 1,000 acres in 1950 to 830 acres in 2001; and during that time about 300 acres were developed for pivot. All of the pivot development occurred prior to 1976. Pivot irrigation has encroached into the active river corridor; approximately 27 acres of pivot-irrigated land is within the natural Channel Migration Zone (CMZ) of the river, making it especially susceptible to threats of river erosion. This pivot is at RM 190R, where a ~300 acre pivot field extends to within 150 feet of the river bank.

Reach C16 shows an increase in forest area considered to be at low risk of cowbird parasitism. In 1950, the reach had 54.5 acres of such forest per valley mile and by 2001 that forest extent had increased to 66.7 acres per valley mile.

A total of 170 acres of Russian olive were mapped in the reach, which is an abrupt increase relative to the two reaches upstream. The Russian olive is distributed throughout the riparian corridor but becomes more prolific in the downstream direction towards Miles City.

Reach C16 was sampled as part of the fisheries study. A total of 32 fish species were sampled in the reach, including Blue Sucker and Sauger, which have been identified as Species of Concern (SOC) by the Montana Natural Heritage Program.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The 100-year flood has dropped by 18 percent and the 2-year flood, which strongly influences overall channel form, has dropped by 24 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 4,850 cfs to 3,070 cfs with human development, a reduction of 37 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 6,340 cfs under unregulated conditions to 3,390 cfs under regulated conditions, a reduction of 47 percent.

Fall and winter base flows have increased in Reach C16 by about 60 percent.

CEA-Related observations in Reach C16 include:

- Pivot irrigation encroachment into CMZ

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach C16 include:

- Russian olive removal
- Removal of flanked rock riprap at RM 190.5R to prevent accelerated erosion behind

Yellowstone River Reach Narratives

Reach C16


The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

Discharge		Undev.	Developed	% Change	"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.		
2 Year (cfs)		62,000	47,300	-23.7%			
100 Year (cfs)		120,000	98,500	-17.9%			
Bankfull Channel Area (Ac)		1950	1976	1995	2001		
		848.9	841.5	827.6	839.3		
					-9.6		
Physical Features	2011 Length (ft)	% of Bankline	2001-2011 Change	There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.			
Rock RipRap	7,009	9.2%	221				
Concrete Riprap	2,192	2.9%	0				
Flow Deflectors	1,555	2.0%	-55				
Total	10,756	14.1%	166				
Length of Side Channels Blocked (ft)	Pre-1950s	Post-1950s	Numerous side channels have been blocked by small dikes.				
	0	0					
Floodplain Turnover	1950 - 1976	1976 - 2001	1950-2001 In-channel riparian encroachment (negative number indicates retreat)		The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.		
Total Acres	120.7	119.2					
Acres/Year	4.6	4.8					
Acres/Year/Valley Mile	0.7	0.7	54.51 acres				
Open Bar Area	Point Bars	Bank Attached	Mid-Channel	Total	The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.		
Change in Area '50 - '01 (Ac)	10.5	46.1	-3	53.6			
Floodplain Isolation	Acres	% of FP	Floodplain isolation refers to area that historically was flooded, but has become isolated due to flow alterations or physical features such as levees.				
5 Year	721.5	62%					
100 Year	308.2	13%					
Restricted Migration Area	Acres	% of CMZ	Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.				
	210.4	9%					
Land Use	1950	2011	1950	2011	Changes in land use reflect the development of the river corridor through time. The irrigated agricultural area is a sub-set of the mapped agricultural land.		
Agricultural Land (Ac)	6,183.9	6,007.7	Flood (Ac)	1,003.6	827.0		
Ag. Infrastructure (Ac)	91.9	159.1	Sprinkler (Ac)	0.0	0.0		
Exurban (Ac)	74.5	3.7	Pivot (Ac)	0.0	303.6		
Urban (Ac)	108.3	366.0					
Transportation (Ac)	117.5	90.6					
1950s Riparian Vegetation Converted to a Developed Land Use (ac)	To Irrigated	To Other Use	Total Rip. Converted	% of 1950s Rip.	Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.		
	1.2	8.3	9.5	1.0%			
National Wetlands Inventory	Acres	Acres per Valley Mi	Total Wetland Acres	Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).			
Riverine	21.2	3.2	139.1				
Emergent	94.7	14.3					
Scrub/Shrub	23.1	3.5					
Russian Olive (2001) (Appx. 100-yr Floodplain)	Acres	%	Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.				
	170.2	3.7%					
Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)	1950	1976	2001	Change 1950-2011	Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.		
	54.5	53.7	66.7	12.2			

Yellowstone River Reach Narratives

Reach C16

PHYSICAL FEATURES MAP (2011)



Yellowstone River Reach Narratives

Reach C16

CHANNEL MIGRATION ZONE MAP

