

County	Treasure	Upstream River Mile	275
Classification	UA: Unconfined anabranching	Downstream River Mile	269.4
General Location	Mission Valley	Length	5.60 mi (9.01 km)

### Narrative Summary

Reach C6 is located in the Mission Valley north of Hysham. The reach is a 5.6 mile long Unconfined Anabranching reach type, indicating minimal valley wall influence and extensive side channels and forested islands. In this area the alluvial valley bottom is approximately 2.5 miles wide, and this broad valley has formed in the relatively erodible Cretaceous-age Bearpaw Shale.

There are just over 3,000 feet of bank armor in the reach, which covers 5.1 percent of the total bankline. About 600 feet of a floodplain dike at RM 273.2R appears to have been eroded out since 2001.

Almost 11,000 feet of side channels have been blocked by physical features in the reach since 1950. One floodplain dike that blocked a side channel at RM 227.8L in 2001 was eroded out and has since been rebuilt. Additional side channel length has been lost passively, overall, there has been about a three mile reduction in side channel length in this reach since 1950.

About 20 percent of the total 100-year floodplain has become isolated due to human development. The 5-year floodplain is even more affected; 70 percent of the historic 5-year floodplain is no longer inundated at that frequency. The isolation of the historic 5-year floodplain, due primarily to flow alterations, has been associated with increased development in these areas; currently there are about 650 acres of flood irrigated land and 200 acres of pivot land within the historic 5-year floodplain. The vast majority of isolated 5-year floodplain area is within irrigated fields south of the river, and the isolation appears to be due to both flow alterations and agricultural dikes.

Land use is dominated by agriculture, with 188 acres of pivot irrigation development since 1950. There are about 260 acres of flood irrigated land within the CMZ, but due to the lack of bank armor, none of the CMZ has become restricted.

Riparian mapping data show a net gain of 158 acres of woody vegetation into the active channel corridor since 1950. This has occurred both on migrating point bars that have become vegetated, as well as within abandoned side channels. Since 1950, the total area of open timber increased by approximately 250 acres. There are 40 acres of Russian olive in the reach.

Reach C6 was sampled as part of the fisheries study. A total of 26 fish species were sampled in the reach.

Reach C6 was sampled as part of the avian study. A total of 32 bird species were identified in the reach. Two bird species identified by the Montana Natural Heritage Program as Potential Species of Concern (PSOC) were found, the Ovenbird, and the Chimney Swift. In contrast to most reaches, Reach C6 has seen an increase in the forested area that is at low risk of cowbird parasitism since 1950. At that time, there were 55 acres per valley mile of such forest, and that number increased to 106 acres per valley mile by 2001.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The 2-year flood, which strongly influences overall channel form, has dropped by 23 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 4,630 cfs to 2,960 cfs with human development, a reduction of 36 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 6,150 cfs under unregulated conditions to 3,320 cfs under regulated conditions at Reach C10 downstream where the analysis begins, a reduction of 46 percent.

CEA-Related observations in Reach C6 include:

- Active and passive loss of thousands of feet of side channel
- Reconstruction of side-channel blockage following its failure post-2001.


Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach C6 include:

- Side channel reactivation at RM 275R and RM 271L
- Russian olive removal

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

<b>Discharge</b>	<b>Undev.</b>	<b>Developed</b>	<b>% Change</b>	<b>"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.</b>		
2 Year (cfs)	61,000	47,000	-23.0%			
100 Year (cfs)	120,000	100,000	-16.7%			
<b>Bankfull Channel Area (Ac)</b>	<b>1950</b>	<b>1976</b>	<b>1995</b>	<b>2001</b>	<b>1950-2001</b>	<b>Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood.</b>
	654.7	611.0	545.8	548.9	-105.8	
<b>Physical Features</b>	<b>2011 Length (ft)</b>	<b>% of Bankline</b>	<b>2001-2011 Change</b>	<b>There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.</b>		
Rock RipRap	2,478	4.1%	0			
Concrete Riprap	574	1.0%	0			
Flow Deflectors	0	0.0%	0			
<b>Total</b>	<b>3,052</b>	<b>5.1%</b>	<b>0</b>			
<b>Length of Side Channels Blocked (ft)</b>	<b>Pre-1950s</b>	<b>Post-1950s</b>	<b>Numerous side channels have been blocked by small dikes.</b>			
	0	10,910				
<b>Floodplain Turnover</b>	<b>1950 - 1976</b>	<b>1976 - 2001</b>	<b>1950-2001 In-channel riparian encroachment (negative number indicates retreat)</b>		<b>The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.</b>	
Total Acres	123.2	92.5	158.33 acres			
Acres/Year	4.7	3.7				
Acres/Year/Valley Mile	1.4	1.1				
<b>Open Bar Area</b>	<b>Point Bars</b>	<b>Bank Attached</b>	<b>Mid-Channel</b>	<b>Total</b>	<b>The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.</b>	
Change in Area '50 - '01 (Ac)	-9.2	7.6	0.3	-1.4		
<b>Floodplain Isolation</b>	<b>Acres</b>	<b>% of FP</b>	<b>Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.</b>			
5 Year	1,663.9	70%				
100 Year	731.8	20%				
<b>Restricted Migration Area</b>	<b>Acres</b>	<b>% of CMZ</b>	<b>Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.</b>			
	176.0	8%				
<b>Land Use</b>	<b>1950</b>	<b>2011</b>	<b>1950</b>	<b>2011</b>	<b>Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land.</b>	
Agricultural Land (Ac)	3,400.5	3,584.1	Flood (Ac)	1,754.0	1,365.9	
Ag. Infrastructure (Ac)	34.4	48.3	Sprinkler (Ac)	0.0	0.0	
Exurban (Ac)	0.0	0.0	Pivot (Ac)	0.0	187.6	
Urban (Ac)	0.0	0.0				
Transportation (Ac)	16.0	16.6				
<b>1950s Riparian Vegetation Converted to a Developed Land Use (ac)</b>	<b>To Irrigated</b>	<b>To Other Use</b>	<b>Total Rip. Converted</b>	<b>% of 1950s Rip.</b>	<b>Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.</b>	
	5.9	0.0	5.9	1.0%		
<b>National Wetlands Inventory</b>	<b>Acres</b>	<b>Acres per Valley Mi</b>	<b>Total Wetland Acres</b>	<b>Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).</b>		
Riverine	19.0	5.5	<b>130.5</b>			
Emergent	89.1	25.8				
Scrub/Shrub	22.5	6.5				
<b>Russian Olive (2001) (Appx. 100-yr Floodplain)</b>	<b>Acres</b>	<b>%</b>	<b>Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.</b>			
	40.0	0.9%				
<b>Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)</b>	<b>1950</b>	<b>1976</b>	<b>2001</b>	<b>Change 1950-2011</b>	<b>Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.</b>	
	54.8	86.2	106.1	51.3		

## PHYSICAL FEATURES MAP (2011)



## CHANNEL MIGRATION ZONE MAP

