

County	Dawson	Upstream River Mile	89
Classification	PCA: Partially confined anabranching	Downstream River Mile	81.4
General Location	Downstream of Glendive	Length	7.60 mi (12.23 km)

Narrative Summary

Reach D7 is located just downstream of Glendive. It is 7.6 miles long and is a Partially Confined Anabranching (PCA) reach type, including some valley wall influence as well as numerous forested islands. These reach types tend to be relatively dynamic with high rates of channel change through time. The Stipek Fishing Access Site is located in the middle portion of the reach.

No bank armor has been mapped in Reach D7, and no side channels have been blocked by dikes. About two miles of transportation encroachment by the railroad was mapped in Reach D7, all of which was in place by 1950.

Similar to many reaches in the Lower Yellowstone Valley, the river channel in Reach D7 has gotten smaller since 1950. The channel contracted by about 121 acres in this reach since 1950, and about 150 acres of riparian vegetation has encroached into old channel areas. This pattern has been consistent in the lower river, and relates primarily to a reduction in flows due to human development. Floodplain turnover rates have dropped from 8.9 acres per year pre-1976 to 5.4 acres per year post-1976.

Even though no side channels have been intentionally blocked, Reach D7 has lost about 3,800 feet of side channel length since 1950. This is likely due to passive loss caused by a reduction in high flows. Lower flows have also resulted in the isolation of 48 percent of the historic 5-year floodplain.

Land use is predominantly agricultural, with about 258 acres of pivot irrigation development since 1950. There are 27 acres of pivot irrigation and 21 acres of exurban land uses in the Channel Migration Zone. Two dump sites have been mapped on the right bank at RM 84R and RM 85.9R.

There are 7.4 acres of mapped Russian olive in the reach.

Reach D7 was part of the avian study. A total of 43 species were identified in the reach, including the Ovenbird, which has been identified by the Montana Natural Heritage Program as a Potential Special Concern. The Black-billed Cuckoo and Red-headed Woodpecker were also identified, both of which are Species of Concern.

A hydrologic evaluation of flow depletions indicates that flow alterations over the last century have been major in this reach. The magnitude of the 100-year flood is now 127,000 cfs, which is 12 percent lower than it was pre-development (145,000 cfs). The 2-year flood, which strongly influences overall channel form, has dropped by 22 percent. Low flows have also been impacted; severe low flows described as 7Q10 (the lowest average 7-day flow anticipated every ten years) for summer months has dropped from an estimated 4,700 cfs to 2,600 cfs with human development, a reduction of 45 percent. More typical summer low flows, described as the summer 95% flow duration, have dropped from 6,890 cfs under unregulated conditions to 3,110 cfs under regulated conditions, a reduction of 55 percent.

Seasonal low flows have increased by 78 percent in the winter and 62 percent in the fall. Both fall and winter base flows are currently about 3,500 cfs.

CEA-Related observations in Reach D7 include:

- Passive loss of side channels with flow alterations

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach D7 include:

- Russian olive removal

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

Discharge	Undev.	Developed	% Change	"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.		
2 Year (cfs)	69,500	54,200	-22.0%			
100 Year (cfs)	145,000	127,000	-12.4%			
Bankfull Channel Area (Ac)	1950	1976	1995	2001	1950-2001	Bankfull channel area is the total footprint of the river inundated at approx. the 2-year flood.
	1,223.9	1,230.6	1,141.1	1,102.9	-121.1	
Physical Features	2011 Length (ft)	% of Bankline	2001-2011 Change	There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.		
Rock RipRap	0	0.0%	0			
Concrete Riprap	0	0.0%	0			
Flow Deflectors	0	0.0%	0			
Total	0	0.0%	0			
Length of Side Channels Blocked (ft)	Pre-1950s	Post-1950s	Numerous side channels have been blocked by small dikes.			
	0	0				
Floodplain Turnover	1950 - 1976	1976 - 2001	1950-2001 In-channel riparian encroachment (negative number indicates retreat)		The rate of floodplain turnover reflects how many acres of land are eroded by the river. Turnover is associated with the creation of riparian habitat.	
Total Acres	230.7	133.9	149.38 acres			
Acres/Year	8.9	5.4				
Acres/Year/Valley Mile	1.3	0.8				
Open Bar Area	Point Bars	Bank Attached	Mid-Channel	Total	The type and extent of open sand and gravel bars reflect in-stream habitat conditions that can be important to fish, amphibians, and ground-nesting birds such as least terns.	
Change in Area '50 - '01 (Ac)	-52.3	40.4	-2.8	-14.6		
Floodplain Isolation	Acres	% of FP	Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.			
5 Year	395.2	48%				
100 Year	43.6	2%				
Restricted Migration Area	Acres	% of CMZ	Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.			
	6.0	0%				
Land Use	1950	2011	1950	2011	Changes in land use reflect the development of the river corridor through time. The irrigated agricultural are is a sub-set of the mapped agricultural land.	
Agricultural Land (Ac)	4,756.4	4,620.5	Flood (Ac)	0.0	708.1	
Ag. Infrastructure (Ac)	29.3	83.7	Sprinkler (Ac)	0.0	25.5	
Exurban (Ac)	0.0	48.9	Pivot (Ac)	0.0	258.3	
Urban (Ac)	0.0	0.0				
Transportation (Ac)	88.2	90.2				
1950s Riparian Vegetation Converted to a Developed Land Use (ac)	To Irrigated	To Other Use	Total Rip. Converted	% of 1950s Rip.	Changes in the extents of riparian vegetation are influenced by land use changes within the corridor.	
	57.6	19.8	77.4	5.0%		
National Wetlands Inventory	Acres	Acres per Valley Mi	Total Wetland Acres	Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), Emergent (marshes and wet meadows) and Shrub-Scrub (open bar areas with colonizing woody vegetation).		
Riverine	28.9	4.2	148.2			
Emergent	72.3	10.6				
Scrub/Shrub	47.1	6.9				
Russian Olive (2001) (Appx. 100-yr Floodplain)	Acres	%	Russian olive is considered an invasive species and its presence in the corridor is fairly recent. Its spread can be used as a general indicator of invasive plants within the corridor.			
	7.4	0.2%				
Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)	1950	1976	2001	Change 1950-2011	Cowbirds are associated with agricultural and residential development, displacing native bird species by parasitizing their nests.	
	116.2	85.5	108.4	-7.9		

CHANNEL MIGRATION ZONE MAP

