Reach D15

County Classification General Location Mckenzie PCM/I: Partially confined meandering/islands Downstream of Fairview Upstream River Mile13.5Downstream River Mile7.5Length6.00 mi (9.66 km)

Narrative Summary

Reach D15 is located downstream of Fairview. The reach is a 6 mile long Partially Confined Meandering with Islands (PCM/I), indicating some valley wall influence, and a meandering main thread with cutoff channels through meander cores forming persistent forested islands.

No bank armor was mapped in the reach, and no side channels have been blocked.

Similar to many reaches in the Lower Yellowstone Valley, the river channel in Reach D15 has gotten smaller since 1950. The channel contracted by about 190 acres in this reach since 1950, and about 210 acres of riparian vegetation has encroached into old channel areas. This pattern has been consistent in the lower river, and relates primarily to a reduction in flows due to human development.

Land use is predominantly agricultural, with 71 acres of pivot irrigation development since 1950. A total of 54 percent of the 100 year floodplain has become isolated (1,885 acres), and most of this isolation is from agricultural dikes. Approximately 23 percent of the 5-year floodplain has become isolated (168 acres).

There is a drill pad on the edge of the CMZ at RM 10.8L.

One ice jam was reported in the reach. It was a break-up flood event on February 12, 1996.

Reach D15 was sampled as part of the avian study. A total of 30 bird species were identified in the reach. Two bird species identified by the Montana Natural Heritage Program as Potential Species of Concern (PSOC) on the Yellowstone River were found, the Ovenbird and the Plumbeous Vireo.. Reach D15 has seen a decrease in the forested area that is at low risk of cowbird parasitism since 1950. At that time, there were 25.6 acres per valley mile of such forest, and that number dropped to 19.6 acres per valley mile by 2001.

CEA-Related observations in Reach D15 include:

•Flow alteration impacts on floodplain access

Recommended Practices (may include Yellowstone River Recommended Practices--YRRPs) for Reach D15 include: • Russian olive removal

The following table summarizes some key CEA results that have been used to describe overall condition and types of human influences affecting the river. The values are specific to this single reach. Blanks indicate that a particular value was not available for this area. This information is consolidated from a large dataset that is presented in more detail in the full reach narrative report.

Discharge 2 Year (cfs) 100 Year (cfs)	Undev. 69,900 143,000	Developed 54,300 134,000	% Change -22.3% -6.3%	"Undeveloped" flows represent conditions prior to significant human development, whereas "developed" flows reflect the current condition of both consumptive and non-consumptive water use.				
Bankfull Channel Area (Ac)	1950 988.3	1976	1995 887.9	2001 798.9	1950-20 -189.3		ful channel area is the total footprint of the inundated at approx. the 2-year flood.	
Physical Features Rock RipRap Concrete Riprap Flow Deflectors	2011 Length (ft) 0 0 0	% of Bankline 0.0% 0.0% 0.0%	2001-2011 Change 0 0 0	There are additional types of bank armor such as car bodies and steel retaining walls, but they are relatively minor.				
Total	0	0.0%	0					
Length of Side Channels Blocked (ft)	Pre-1950s 0	Post-1950s 0		Numerous side channels have been blocked by small dikes.				
Floodplain Turnover Total Acres Acres/Year Acres/Year/Valley Mile	1950 - 1976	1976 - 2001	rip	1950-2001 In-channel riparian encroachment (negative number indicates retreat) 208.49 acresThe rate of floodplain turnover reflects how many acres of land are eroded by the river. Tunover is associated with the creation of riparian habitat.				
Open Bar Area Change in Area '50 - '01 (Ac)	Point Bars 0	Bank Attached 89.7	Mid- Channel -57.5	The type and extent of open sand and gravel bars reflect in-Totalstream habitat conditions that can be important to fish,32.2amphibians, and ground-nesting birds such as least terns.				
Floodplain Isolation 5 Year 100 Year	Acres 168.1 1,884.7	<mark>% of FP</mark> 23% 54%		Floodplain isolation refers to area that historically was flooded, but has become isolated do to flow alterations or physical features such as levees.				
Restricted Migration Area	Acres 21.1	<mark>% of CMZ</mark> 1%	Channel Migration Zone restrictions refer to the area and percent of the CMZ that has been isolated by features such as bank armor, dikes, levees, and transportation embankments.					
and Use	1950	2011			1950	2011	Changes in land use reflect the	
Agricultural Land (Ac)		7,485.3	Flood (A	Ac)	3,955.0	6,101.5	development of the river corridor through	
Ag. Infrastructure (Ac)	86.2	192.8	Sprinkle	er (Ac)	0.0	0.0	time. The irrigated agricultural are is a sub-set of the mapped agricultural land.	
Exurban (Ac) Urban (Ac)	0.0 0.0	35.8 0.0	Pivot (A		0.0	71.3		
Transportation (Ac)	79.3	70.6						
L950s Riparian Vegetation Converted to a Developed .and Use (ac)	To Irrigated	To Other Use	Total Rip. Converted	% of 1950s Rip.	changes in the extents of upartan vegetation are innacheed by			
National Wetlands Inventory	Acres	Acres per Valley Mi	Тс	Wetlands units summarized from National Wetlands Inventory Mapping include Riverine (typically open water sloughs), and Emergent (marshes and wet meadows) and Shrub-Scrub (open res bar areas with colonizing woody vegetation)5				
Riverine Emergent Scrub/Shrub	1.6 20.2 68.7	0.3 3.5 11.9						
Russian Olive (2001) Appx. 100-yr Floodplain)	Acres 0.8	<mark>%</mark> 0.1%		s considered an invasive species and its presence in the corridor is fairly recent. be used as a general indicator of invasive plants within the corridor.				
Riparian Forest at low risk of Cowbird Parasitism (Ac/Valley Mile)	1950 10.1	1976	2001 23.0	Change 1950-2011 12.9			ated with agricultural and residential acing native bird species by parasitizing their	

Reach D15

PHYSICAL FEATURES MAP (2011)



Reach D15

CHANNEL MIGRATION ZONE MAP

